

Table 2: Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
Asia Pacific, High Income					
	Japan	Shingu ¹	1990–1992	40.2 (Estimated number 14,968; Males, n=7,842, Females, n=1,909)	Land transport 44%, n=4,263, Falls 42%, n=4,078, Sports/recreation 5%, n=528, Self-harm 2%, n=167 and Other/unknown 7%, n=716
	Japan (Okayama Prefecture)	Ide ²	1988-1989	28.6 (n=55; Males, n=45, Females, n=10)	Land transport 33%, n=18, Falls 16%, n=9, Work-related 38%, n=21 and Other/unknown 13%, n=7
Asia, East					
	China	Zhao ³	1983 and 1985	n/r*	Land transport 12.1%-16.7%, Falls 33.3%-66.7% and Violence 15.1%-22.2%
	Taiwan	Yeh ⁴	1977-1989	250 cases annually** (total cases n=1,617; Males, n=1,295, Females n=322)	Land transport 63%, n=1,096, Falls and Self-harm 12%, n=200, Work-related 19%, n=308 and Other/unknown 6%, n=13
	Taiwan	Chen ⁵	1992–1996	18.8 (n=1,232; Males, n=958, Females, n=274)	Land transport 49%, n=609, Falls 41%, n=500, Sports/recreation 1%, n=16, Violence 2%, n=25 and Other/unknown 7% n=82 (Age group 15-64 years)
	Taiwan	Chen ⁵	1992–1996	47.5 (Geriatric population, n=297; Males, n=186, Females, n=111)	Land transport 33%, n=97, Falls 62%, n=185, Sports/recreation 0.3%, n=1, Violence 2%, n=5 and Other/unknown 2.7%, n=9 (Age group 65-and above years)
	Taiwan	Yang ⁶	2000-2003	73.0 (n=6,455); Males, n=3,058, Females*, n=3,397) *ICD9 806.X (Fractured vertebra with SCI and assuming male to female ratio 0.9	n/r*
	Taiwan (Taipei)	Chen ⁷	1978-1981	14.6 (n=560; Males, n=465, Females, n=95)	Land transport 45%, n=249, Falls 29%, n=157, Sports/recreation 3%, n=19, Violence 3%, n=15, Suicide 1%, n=8 and Other/unknown 20%, n=112

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
Asia, South	Taiwan (Hualien County)	Lan ⁸	1986-1990	56.1(n=99; Males, n=79, Females, n=20)	Land transport 62%, n=61, Falls 23%, n=23, Sports/recreation 6%, n=6, Violence 3%, n=3, Self-harm 2%, n=2 and Other/unknown 4%, n=4
	Bangladesh	Hoque ⁹	1994-1995	n/r (n=179; Males, n=158, Females, n=21)	Land transport 18%, n=33, Falls 63%, n=113 and Other/unknown 19%, n=33
	Nepal	Mukhida ¹⁰	2001-2004	5 cases annually (Estimate, ages <19 years)(n=14 during study period; Males, n=9, Females, n=5)	Falls 61%, n=9
	Nepal (East)	Lakhey ¹¹	1997-2001	n/r (n=233; Males, n=169, Females, n=64)	Land transport 7%, n=16, Falls 82%, n=192 and Other/unknown 11%, n=25
	Pakistan	Raja ¹²	1995-1999	5.1 (n=2,654; Males, n=1,922, Females, n=732)	Land transport 7%, n=186, Falls 82%, n=2,176 and Other/unknown 11%, n=292
	Pakistan	Masood ¹³	2003-2007	n/r (n=214; Males, n=189, Females, n=25)	Land transport 23%, n=50, Falls 68%, n=145, Sports/recreation 1%, n=3, Violence 6%, n=12 and Other/unknown 2%, n=4
	Pakistan (AFIRM, Rawalpindi)	Rathore ¹⁴	2006	n/r (n=83; Males, n=68, Females 15)	Land transport 25%, n=21, Falls 58%, n=48, Violence 8%, n=7 and Other/unknown 9%, n=7
Asia, Southeast	Thailand (Chiang Mai)	Kovindha ¹⁵	1985-1991	23 (n=398; Males, n=365, Females, n=33)	Land transport 47%, n=187, Falls 48%, n=191, Violence 4%, n=16 and Other/unknown 1%, n=4
	Thailand (Bangkok)	Parajareya ¹⁶	1989-1994	5.8 (Estimate based on 1992 population of 6,250,000; n=219; Males, n=186, Females, n=33)	Land transport 51%, n=111, Falls 31%, n=67, Sport/recreation 0.5%, n=1, Violence 8%, n=18, Self-harm 0.5%, n=1 and Other/unknown 9%, n=21
	Viet Nam	Weerts ^{17, 18}	2006-2007	13.9 (Estimate based on 2008 population of 86,116,560; n=247)	Land transport 45%, n=111, Sports/recreation 2%, n=4, Work-related 17%, n=40, Violence 1%, n=3 and Other/unknown 36%, n=89

Continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
Australasia					
	Australia	O'Connor ¹⁹	1998-1999	14.5 (Age-standardised, n=265; Males, n=201, Females, n=64)	Land transport 43%, n=113, Falls 31%, n=83, Sports/recreation 9%, n=24 and Other/unknown 17%, n=45
	Australia	Cripps ²⁰	2006-2007	14.9 (Age-standardised, n=272; Males, n=226, Females, n=46)	Land transport 52%, n=142, Falls 29%, n=78, Sports/recreation 8%, n=21 and Other/unknown 11%, n=30
	Australia (Victoria)	Cheshire ²¹	1959-1966	11.6 (Based on Victorian population of 4 million; n=390; Males, n=328, Females, n=62)	Land transport 52%, n=170, Sports/recreation 13%, n=41, Work-related 21%, n=68, Violence 2%, n=5 and Other/unknown 12%, n=41
	Australia (Queensland)	Sutton ²²	1962-1971	16.6 (cases >9 years of age; estimated incidence based on 1.75 million population, n=232; Males, n=202, Females, n=30)	Land transport 50%, n=103, Falls 7%, n=15, Sports/recreation 21%, n=37, Work-related 20%, n=40 and Violence 2%, n=12
	Australia (Victoria)	Burke ²³	1978-1982	325 cases during reporting period; Males, n=254, Females, n=71	Land transport 52%, n=169, Sports/recreation 16%, n=52, Work-related 12%, n=40 and Other/unknown 20%, n=65
	New Zealand	Dixon ²⁴	1988	49.1 (n=164; Males, n=120, Females, n=44)	Land Transport 54%, n=89, Falls 24%, n=39, Sports/recreation 11%, n=18, Violence 2%, n=3 and Other/unknown 9%, n=15
Europe, Central					
	Poland (Poznan)	Jankowski ²⁵	1979-1992	n/r (n=511; Males, n=449, Females, n=62)	Land transport 41%, n=208, Sports/recreation 19%, n=97, Work-related 15%, n=79 and Other/unknown 25%, n=127
	Romania (Bucharest)	Soopramanien ²⁶	1975-1992	28.5 (n=400/yr approx; Males, n=317, Females, n=95)	Land transport 72%, n=296, Sports/recreation 7%, n=29, Work-related 13%, n=52 and Other/unknown 8%, n=35

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
Europe, Eastern	Estonia	Sabre ²⁷	2003-2007	27.9 (n=191; Males, n=159, Females, n=32)	Land transport 27%, n=52, Falls 42%, n=80, Sports/recreation 15%, n=29, Self-harm 1%, n=2, Violence 4%, n=8 and Other/unknown 11%, n=20
	Russia (Novosibirsk)	Silberstein ²⁸	1989-1993	29.7 (n=196 over period 5 yrs; Males, n=153, Females, n=43)	Land transport 25%, n=49, Falls 40%, n=79, Sports/recreation 33%, n=65 and Violence 2%, n=3
	Russia (Saint Petersburg)	Kondakov ²⁹	1994-1996	44 (n=105; Males, n=68, Females, n=37).	Land transport 16%, n=17, Falls 34%, n=36, Sports/recreation 2%, n=2, Work-related 5%, n=5 and Other/unknown 43%, n=45
Europe, Western	Denmark	Biering-Sorensen ³⁰	1975-1984	9.2 (n=360 over this period; Males, n=262, Females, n=98)	Land transport 47%, n=125, Falls 26%, n=70, Sports/recreation 12%, n=30, Self-harm 8%, n=21, Violence 2%, n=5 and Other/unknown 5%, n=17
	Finland	Kannus ³¹	1970-1995	50 (Fall-related SCI in 50 years and above in 1970); 270 (Fall-related SCI in 50 years and above in 1995)	n/r
	Finland (Helsinki)	Dahlberg ³²	1978-1997	13-15 (n=55 cases per year; Males, n=42, Females, n=13)	Land transport 35%, n=53, Falls 43%, n=66, Sports/recreation 9%, n=14, Violence 4%, n=6 and Other/unknown 9%, n=13
	Finland	Ahoniemi ³³	1976-2005	13.8 (n=1,647 >16yrs; Males, n=1,362, Females, n=285)	Land transport 40%, n=651, Falls 41%, n=679, Sports/recreation 7%, n=109, Violence 3%, n=44 and Other/unknown 9%, n=164
	France	Albert ³⁴	2000	19.4 (n=934 Ages>15 years)	n/r
	France (Rhone-Alpes Region)	Minaire ¹⁰	1970-1975	12.7 (n=351; Males, n=276, Females, n=75)	Land transport 47%, n=166, Falls 29%, n=100, Sport/recreation 8%, n=27, Violence 1%, n=3 and Other/unknown/ 15%, n=55

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	Germany	Exner ³⁵	1976-1996	10.7 (Based on 1985 pop of 77,685,000; n=16,559 over the reporting period; Males, n=11,827, Females, n=4,732) ^{***}	Land transport 46%, n=7,699, Sports/recreation 10%, n=1,709, Work-related 19%, n=3,119, Self-harm 7%, n=1,080, Violence 1%, n=239 and Other/unknown 17%, n=2,713
	Germany	Koning ³⁶	1983	36 (n=2,213)	n/r
	Greece (Thessaloniki)	Divanoglou ³⁷	2006-2007	33.6 (n=87 >15 yrs; Males, n=77, Females, n=10)	Land transport 51%, n=41, Falls 37%, n=30, Sports/recreation 4%, n=3, Violence 2%, n=2 and Other/unknown 6%, n=5
	Greenland	Pedersen ³⁸	1965-1986	26 (n=27; Males, n=20, Females, n=7)	Land transport 4%, n=1, Falls 33%, n=9, Sports/recreation 4%, n=1, Self-harm 26%, n=7, Violence 11%, n=3, and Other/unknown 22%, n=6
	Iceland	Knuttdottir ³⁹	1973-1989	24 (1973-82); 18 (1983-89) (n=79); Males, n=43, Females, n=16, were living)	Land transport 50%, n=39, Falls 37%, n=29 and Other/unknown 13%, n=11
	Iceland	Knuttdottir ⁴⁰	1973-2008	20 (n=191; Males, n=140, Females, n=51)	Land transport 46%, n=87, Falls 29%, n=55, Sports/recreation 20%, n=39, Other/unknown 5%, n=10.
	Ireland	O'Connor ⁴¹	2000	13.1 (n=46; Males, n=40, Females, n=6)	Land transport 50%, n=23, Falls 22%, n=10, Work-related 15%, n=7, Sports/recreation 9%, n=4, Self-harm 2%, n=1 and Other/unknown 2%, n=1
	Ireland (West Ireland)	Roche ⁴²	4-year study	195.4 * (n=285; Males, n=175, Females, n=111)	Land transport 34%, n=97, Falls 58%, n=168, Sports/recreation 2%, n=4 and Other/unknown 5%, n=16
	Israel	Catz ⁴³	1959-1992	15.9 (n=250 consecutive patients; Males, n=189, Females, n=61) (Based on 1989 pop) ^{***}	Land transport 33%, n=82, Falls 17%, n=42, Work-related 27%, n=67, Self-harm 14%, n=34 and Other/unknown 9%, n=25
	Israel	Ohry ⁴⁴	1974-1984	1 (Approx., paediatric SCI. n=23; Males, n=15, Females, n=8)	Land transport 35%, n=8, Falls 22%, n=5, Sports/recreation 22%, n=5 and Violence 21%, n=5

*Likely inaccurate

Continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	Italy (7 Rehabilitation Centres)	Celani ⁴⁵	1989-1994	2.3 (n=642; Males, n=509, Females, n=133) (Based on 1990 population)***	Land transport 58%, n=375, Sports/recreation 8%, n=27, Work-related 15%, n=98, Violence 3%, n=19, Self-harm 4%, n=27 and Other/unknown 15%, n=96
	Italy	Pagliacci ⁴⁶	1997-1999	Estimated to be 18-20 (n=684; Males, n=548, Females, n=136)	Land transport 54%, n=368, Falls 23%, n=155, Sports/recreation 8%, n=54, Self-harm 4%, n=29 and Other/unknown 11%, n=78
	Italy (Veneto)	Caldana ⁴⁷	1994-1995	14.3 (n=127; Males, n=106, Females, n=21)	Land transport 57%, n=72, Falls 10%, n=13, Work-related 22%, n=28, Sports/recreation 6%, n=8, Violence 1%, n=1 and Other/unknown 4%, n=5
	Netherlands	Schonher ⁴⁸	1982-1993	7.7 (n=142; Males, n=109, Females, n=33)	n/r
	Netherlands	van Asbeck ⁴⁹	1994	7.5 (n=113; Males, n=87, Females, n=26)	Land transport 31%, n=35, Falls 49%, n=55, Work-related 4%, n=5, Sports/recreation 9%, n=10 and Other/unknown 7%, n=8
	Netherlands and Belgium (Flanders)	Osterthun ⁵⁰	2002-2007	n/r (n=503; Males, n=371, Females, n=132)	n/r
	Norway	Lidal ⁵¹	1961-1982	4.5 approx (n=387; Males, n=321, Females, n=66)	Land transport 46%, n=178, Falls 32%, n=124, Sports/recreation 11%, n=43, Violence 4%, n=15 and Other/unknown 7%, n=27
	Norway	Gjone ⁵²	1974-1975	16.5 (n=131; Males, n=109, Females, n=22)	n/r
	Norway	Augutis ⁵³	2002-2004	1.5 (Paediatric SCI aged 0-15 yrs, n=42)	n/r
	Norway (Western)	Hagen ⁵⁴	1952-2001	26.3 (Annual incidence in 1997-2001; n=71; Males, n=277, Females, n=59)	Land transport 34%, n=115, Falls 46%, n=153, Sports/recreation 9%, n=29 and Other/unknown 11%, n=39
	Portugal (Coimbra)	Martins ⁵⁵	1989-1992	57.8 (Survivors plus those who died, n=398; Males, n=306, Females, n=92)	Land transport 57%, n=228, Falls 37%, n=149 and Other/unknown 6%, n=21

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	Portugal	Augutis ⁵³	2002-2004	27 (Paediatric SCI aged 0-14 yrs, n=138)	n/r
	Spain	Herruzo-Cabrera ⁵⁶	1984-1985	8.1*** (n=208)	n/r
	Sweden	Augutis ⁵³	2002-2004	4.6 (Paediatric SCI aged 0-15 yrs, n=21)	n/r
	Sweden (Stockholm)	Divanoglou ³⁷	2006-2007	19.6 (n=49; Males, n=36, Females, n=11 [consenting cases])	Land transport 23%, n=11, Falls 38%, n=18, Sports/recreation 17%, n=8, Self-harm 9%, n=4, Assault 2%, n=1 and Other/unknown 11%, n=5
	Sweden	Augutis ⁵⁷	1985-1996	4.6 (Paediatric SCI aged 0-15 yrs, n=92; Males, n=47, Females, n=45)	Land transport 75%, n=69, Falls 9%, n=8, Sports/recreation 13%, n=12, and Other/unknown 3%, n=3
	Switzerland	Gehrig ⁵⁸	1960-1967	15 (n=584; Males, n=461, Females, n=123)	Land transport 36%, n=210, Sports/recreation 19%, n=111, Work-related 35%, n=204, Self-harm 9%, n=53 and Other/unknown 1%, n=6
	UK	Augutis ⁵³	2002-2004	5.2 (Paediatric SCI aged 0-14 yrs, n=180)	n/r
	UK (Midlands, Nth Wales, Mid Wales, South Mersey)	Aung ⁵⁹	1985-1988	7.3 (n=219; Males, n=173, Females, n=46)	Land transport 50%, n=110, Falls 43%, n=93 and Sports/recreation 7%, n=16
Latin America, Tropical	Brazil	Brasil ⁶⁰	1986-1995	n/r (n=164; Males, n=134, Females, n=30)	Land transport 38%, n=62, Falls 26%, n=42, Sports/recreation 7%, n=12, Violence 12%, n=19 and Other/unknown 17%, n=29
	Brazil	Greve ⁶¹	1986-2007	Estimated 17.3*** (n=3,289; Males, n=2,741, Females, n=548)	Land transport 31%, n=1,010, Falls 39%, n=1,289, Sports/recreation 14%, n=474, and Violence 16%, n=516
	Brazil (San Paulo)	Barros ⁶²	1982-1987	10.1 (Estimated using 1981 population of San Paulo)**** (n=428; Males, n=404, Females, n=24)	Land transport 27%, n=115, Falls 22%, n=96, Sports/recreation 8%, n=34, Violence 37%, n=157 and Other/unknown 6%, n=26

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
North Africa/Middle East	Brazil (7 Brazilian capitals)	Campos da Paz ⁶³	1988	n/r (n=108; Males, n=87, Females, n=21)	Land transport 42%, n=45, Falls 15%, n=16, Sports/recreation 9%, n=10, Violence 28%, n=30 and Other/unknown 9%, n=7
	Jordan	Otom ⁶⁴	1998-1993	18 (n=151; Males, n=129, Females, n=22)	Land transport 44%, n=67, Falls 21%, n=32, Sports/recreation 3%, n=4, Violence 28%, n=42, and Other/unknown 4%, n=6
	Saudi Arabia	Alshahri ⁶⁵	2003-2008	1.6 (based on a population of 24 million; n=307 over the 8-year period; Males n=271, Females, n=36)	Land transport 85%, n=261, Falls 9%, n=28, Sports/recreation 1%, n=3 and Violence 5%, n=15
	Turkey	Dincer ⁶⁶	1974-1985	n/r (n=1,694; Males, n=1,282, Females, n=412)	Land transport 35%, n=600, Falls 30%, n=500, Violence 24%, n=406 and Other/unknown 13%, n=188
	Turkey	Karacan ⁶⁷	1992	12.7 (n=581; Males, n=415, Females, n=166)	Land transport 49%, n=284, Falls 37%, n=212, Sports/recreation 1%, n=7, Violence 5%, n=30 and Other/unknown 8%, n=48
	Turkey (SE Anatolia)	Gur ⁶⁸	1990-1999	12.06 (Average incidence) (n=539; Males, n=416, Females, n=123)	Land transport 37%, n=200, Falls 32%, n=172, Violence 23%, n=115, Self-harm 3%, n=15 and Other/unknown 5%, n=27
	Turkey (Istanbul)	Karamehmetoglu ⁶⁹	1992	21.1 (n=152; Males, n=114, Females, n=38)	Land transport 41%, n=62, Falls 43%, n=65, Violence 7%, n=11 and Other/unknown 5%, n=14
	Turkey (Rural areas)	Karamehmetoglu ⁷⁰	1994	16.9 (n=75; Males, n=64, Females, n=11)	Land transport 25%, n=19, Falls 37%, n=28, Violence 31%, n=23 and Other/unknown 7%, n=5
North America, High Income	Canada	Rick Hansen SCI Register ⁷¹	2001-2002	35 (n=1,050; Males, n=840, Females, n=210)	Land transport 47%, n=494, Falls 17%, n=179, Sports/recreation 7%, n=74 and Other/unknown 29%, n=303
	Canada (Toronto)	Tator ⁷²	1947-1973	2.3 (n=351; Males, n=287, Females, n=64)	Land transport 34%, n=119, Falls 10%, n=35, Sports/recreation 15%, n=54, Work-related 30%, n=105 and Other/unknown 11%, n=38

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	Canada (Quebec)	Barss ⁷³	1961-2004	0.64 (Diving related SCI, n=171) Males, n=82, Females 7, based on 89 survivors	Sports/recreation 100%, n=171
	Canada (British Columbia)	Andrews ⁷⁴	1964-1977	0.66 (Paediatric cases, n=15; Males, n=9, Females, n=6)	Land transport 67%, n=10, Violence 13%, n=2 and Other/unknown 20%, n=3
	Canada (Toronto)	Tator ⁷²	1974-1981	3.6 (n=201; Males, n=160, Females, n=41)	Land transport 41%, n=82, Falls 12%, n=24, Sports/recreation 23%, n=46, Work-related 14%, n=28 and Other/unknown 10%, n=21
	Canada (Alberta)	Hamilton ⁷⁵	1975-1988	9.9 (Paediatric cases 17 yrs and below, based on 1988 population of 1.25 million; n=174; Males, n=115, Females, n=59)	Land transport 60%, n=104, Falls 12%, n=21, Sports/recreation 18%, n=31 and Other/unknown 10%, n=18
	Canada (Manitoba)	Hu ⁷⁶	1981-1984	290 (n=944; Males, n=570, Females, n=374)	n/r
	Canada (Ontario)	Pickett ⁷⁷	1994-1999	From 37.2-46.2 age standardised rates (n=2,385; Males, n=1,632, Females, n=753)	Land transport 43%, n=1,021, Falls 43%, n=1,030, Violence 2%, n=52, Self-harm 1%, n=34 and Other/unknown 11%, n=248
	Canada (Alberta)	Dryden ⁷⁸	1997-2000	52.5 (n=450; Males, n=322, Females, n=128)	Land transport 56%, n=254, Falls 19%, n=86, Sports/recreation 11%, n=51, Violence 10%, n=44 and Other/unknown 4%, n=15
	Canada (Southwest Ontario)	Pickett ⁷⁹	1997-2001	40.8 (n=151; Males, n=112, Females, n=39)	Land transport 35%, n=53, Falls 31%, n=47, Sports/recreation 9%, n=14, Work-related 7%, n=10, Violence 5%, n=7, Self-harm 1%, n=2 and Other/unknown 11%, n=18
	USA	Samsa ⁸⁰	1940-1987	n/r (n=5,545; Male veterans)	Land transport 39%, n=2,163, Sports/recreation 13%, n=721, Violence 35%, n=1,941 and Other/unknown 13%, n=720
	USA	Bracken ⁸¹	1970-1977	40.1 (8-year incidence)	n/r

*continued***Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data**

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	USA	Nobunaga ⁸²	1973-1998	n/r (n=25,054; Males, n=20,419, Females, n=4,635)	Land transport 43%, n=10,773, Falls 19%, n=4,710, Sports/recreation 11%, n=2781, Violence 19%, n=4,735 and Other/unknown 8%, n=2,055
	USA	Jackson ⁸³	1973-2003	40 (n=30,532; Males, n=24,426, Females, n=6,106)	Land transport 46%, n=13,923, Falls 20%, n=5,984, Sports/recreation 11%, n=3,267, Violence 18%, n=5,435 and Other/unknown 5%, n=1,923
	USA	Kalsbeek ⁸⁴	1974	50 (estimated n=10,000; Males, n=5,000, Females, n=5,000)	Land transport 33%, n=3,300, Falls 33%, n=3,300 and Other/unknown 34%, n=3,400
	USA	Burney ⁸⁵	1982-1989	n/r (n=2,741; Males, n=2,165, Females, n=576)	Land transport 52%, n=1,439, Falls 20%, n=543, Violence 15%, n=400 and Other/unknown 13%, n=359
	USA	Stover ⁸⁶	1984	30.0 (n=10,000+; Males, n=8,000, Females, n=2,000)	Land transport 48%, n=4,800, Falls 21%, n=2,100, Sports/recreation 14%, n=1,400, Violence 15%, n=1,500 and Other/unknown 2%, n=200
	USA	Harvey ⁸⁷	1988	n/r (n=758; Males, n=538, Females, n=220)	Land transport 48%, n=364, Falls 17%, n=129, Sports/recreation 16%, n=121, Violence 5%, n=38 and Other/unknown 14%, n=106
	USA	Ditunno ⁸⁸	1993	38.8 (Estimated)	Land transport 45%, Falls 22%, Sports/recreation 13%, Violence 16% and Other/unknown 4%
	USA	Lasfargues ⁸⁹	1994	38 (n=11,500 estimated; Males, n=8,015, Females, n=3,485)	n/r
	USA	Vitale ⁹⁰	1997, 2000	19.9 (Paediatric SCI, ages 0-18 years; n=2,909; Males, n=2,088, Females, n=821)	Land transport 56%, n=1,629, Falls 14%, n=407, Sports/recreation 8%, n=233, Violence 9%, n=262 and Other/unknown 13%, n=378
	USA	Ho ⁹¹	2000-2003	n/r	Land transport 50%, Falls 24%, Sports/recreation 9%, Violence 11% and Other/unknown 6%

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	USA	NSCISC ⁹²	2005	40 (n=11,000 approx.; Males, n=8,756, Females, n=2,244)	Land transport 48%, n=5,280, Falls 23%, n=2,530, Sports/recreation 9%, n=990, Violence 14%, n=1,540 and Other/unknown 6%, n=660
	USA (Olmsted County, Minnesota)	Griffin ⁹³	1935-1981	54.8 (age-standardised rate; n=154; Males, n=111, Females, n=43)	Land transport 69%, n=106, Falls 13%, n=20, Sport/recreation 8%, n=12, work-related 7%, n=11 and Violence 3%, n=5
	USA (California)	Kraus ⁹⁴	1970-1971	53.4 (n=619; Males, n=457, Females, n=162)	Land transport 56%, n=347, Falls 20%, n=124, Sports/recreation 7%, n=43, Violence 12%, n=74 and Other/unknown 5%, n=31
	USA (Arkansas)	Acton ⁹⁵	1980-1989	28.5 (age-adjusted incidence; n=644; Males, n=518, Females, n=126)	Land transport 50%, n=319, Falls 13%, n=86, Sports/recreation 11%, n=73, Violence 11%, n=72 and Other/unknown 15%, n=94
	USA (New York)	MMWR ⁹⁶	1982-1988	43 (n=5,384)	n/r
	USA (Hawaii)	Goebert ⁹⁷	1987-1989	23.7 (Based on 1989 population of 1,245,600; n=59; Males, n=50, Females, n=9)	Land transport 38%, n=22, Falls 28%, n=17, Sports/recreation 19%, n=11 and Violence 15%, n=9
	USA (Oklahoma)	Price ⁹⁸	1988-1990	40 (n=376; Males, n=300, Females, n=76)	Land transport 50%, n=189, Falls 20%, n=75, Sports/recreation 13%, n=48, Violence 11%, n=42 and Other/unknown 6%, n=22
	USA (Utah)	Thurman ⁹⁹	1989-1991	43 (47 cases per million, age adjusted to 1980 US pop; n=223; Males, n=169, Females, n=54)	Land transport 49%, n=110, Falls 21%, n=47, Sports/recreation 16%, n=35, Violence 5%, n=12 and Other/unknown 9%, n=19
	USA (Alaska)	Warren ¹⁰⁰	1991-1993	83 (n=139; Males, n=116, Females, n=23)	Land transport 53%, n=73, Falls 22%, n=30, Violence 6%, n=8, Sport/recreation 6%, n=8 and Other/unknown 13%, n=20
	USA (Mississippi)	Surkin ¹⁰¹	1992-1994	77 (n=395; Males, n=316, Females, n=79)	Land transport 58%, n=230, Falls 13%, n=51, Sport/recreation 4%, n=14, Violence 16%, n=62 and Other/unknown 9%, n=38
	USA (Colorado)	Johnson ¹⁰²	1994	45.1-46.3 (n=165-169)	n/r

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	USA (Kentucky, Indiana)	Burke ¹⁰³	1993-1998	27.1 (n=161; Males, n=121, Females, n=40)	Land transport 55%, n=88, Falls 23%, n=37, Violence 14%, n=23 and Other/unknown 8%, n=13
	USA (Rhode Island)	Buechner ¹⁰⁴	1994-1998	56 (n=277; Males, n=185, Females, n=92)	Land transport 29%, n=80, Falls 39%, n=108, Assault 5%, n=14, Self-harm 1%, n=3 and Other/unknown 26%, n=72
	USA (Dade County)	Calancie ¹⁰⁵	1997-2000	n/r (n=229; Males, n=175, Females, n=54)	Land transport 52%, n=118, Falls 20%, n=45, Sports/recreation 11%, n=26, Violence 11%, n=24 and Other/unknown 7%, n=16
Oceania					
	Fiji Islands	Maharaj ¹⁰⁶	1985-1994	10.0 (n=75; Males, n=65, Females, n=10)	Land transport 25%, n=19, Falls 39%, n=29, Sports/recreation 28%, n=21 and Other/unknown 8%, n=6
Sub-Saharan Africa, Southern					
	South Africa (Cape Province)	Key ¹⁰⁷	1963-1967	n/r (n=300; Males, n=270, Females, n=30)	Land transport 34%, n=120, Falls 12%, n=37, Sports/recreation 5%, n=16, Violence 25%, n=76, Other/unknown 24%, n=51
	South Africa (Johannesburg)	Velmahos ¹⁰⁸	1988-1992	48.5 (Median, based on 1990 population of 2,554,726; n=551; Males, n=428, Females, n=123)	Land transport 30%, n=165; Violence 61%, n=336 and Other/unknown 9%, n=50
	South African (Southern Transvaal)	Hart ¹⁰⁹	1988-1993	n/r (n=546; Males, n=437, Females, n=109)	Land transport 28%, n=152, Falls 3%, n=15, Violence 62%, n=341 and Other/unknown 7%, n=38
	Zimbabwe	Levy ¹¹⁰	1988-1994	11.7 (Based on estimated population of 11.6 million; n=136; Males, n=121, Females, n=15)	Land transport 50%, n=68, Falls 17%, n=23, Violence 15%, n=20 and Other/unknown 18%, n=25
Sub-Saharan Africa, West					
	Nigeria (Zaria)	Iwegbu ¹¹¹	1973-1982	n/r (n=48; Males, n=42, Females, n=6)	Land transport 75%, n=36, Falls 8%, n=4 and Other/unknown 17%, n=6
	Nigeria (Plateau State)	Igun ¹¹²	1984-1997	5 cases per year (n=68; Males, n=62, Females, n=6)	Land transport 57%, n=39, Falls 22%, n=15, Work-related 18%, n=12 and Violence 3%, n=2

continued

Table 2 (continued): Incidence and aetiology of traumatic spinal cord injury by Region and Author(s) of published data

Region	Country	Author(s) of published data	Observation period	Incidence per million population per year	Aetiology of SCI
	Nigeria (Lagos)	Obalum ¹¹³	1992-2006	n/r (n=468; Males, n=328, Females, n=140)	Land transport 77%, n=362, Falls 9%, n=44, Sports/recreation 2%, n=8, Violence 11%, n=52 and Other/unknown 1%, n=2
	Nigeria (Ilorin)	Solagberu ¹¹⁴	1995-1999	n/r (n=39; Males, n=36, Females, n=3)	Land transport 67%, n=26, Falls 23%, n=9 and Other/unknown 10%, n=4
	Nigeria (Enugu)	Nwankwo ¹¹⁵	1996-2000	n/r (n=74)	n/r
	Nigeria (Enugu)	Nwadinigwe ¹¹⁶	1996-2001	n/r (n=104; Males, n=88, Females, n=16)	Land transport 58%, n=60, Falls 24%, n=25, Violence 12%, n=12 and Other/unknown 6%, n=7
	Nigeria (Ile-Ife)	Olasode ¹¹⁷	Estimate: 2004-2006	n/r (n=71; Males, n=47, Females, n=24)	Land transport 89%, n=63, Violence 4%, n=3, Self-harm 1%, n=1 and Other/unknown 6%, n=5
	Sierra Leone	Gosselin ¹¹⁸	2002-2004	3.4*** (n=24; Males, n=22, Females, n=2)	Land transport 50%, n=12, Fall 42%, n=10 and Other/unknown 8%, n=2

* n/r: Not reported

** Figures in italics indicate 'Incidence per million population per year' was not available

*** Source/: Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat, /World Population Prospects: The 2008 Revision/, <http://esa.un.org/unpp>

**** Source: http://en.wikipedia.org/wiki/São_Paulo#Population_growth

***** Estimated incidence based on 1986-2007 case counts (3,289) and 2007 Brazilian population of 190,010,647.

1. Shingu H, Ohama M, Ikata T, Katoh S, Akatsu T. A nationwide epidemiological survey of spinal cord injuries in Japan from January 1990 to December 1992. *Paraplegia* 1995; **33**(4): 183-188.
2. Ide M, Ogata H, Tokuhiko A, Takechi H. Spinal Cord Injuries in Okayama Prefecture: An epidemiological Study '88-'89. *J UOEH* 1993; **15**(3): 209-215.
3. Zhao Y-D, Wang W. Neurosurgical Trauma in People's Republic of China. *World J. Surg.* 2001; **25**: 1202-1204.
4. Yeh Y-S, Lee S-T, Liui T-N, Fairholm DJ, Chen W-J, Wong M-K. Features of Spinal Cord Injury in Taiwan (1977-1989). *Chang Gung Med J* 1993; **16**(3): 170-175.
5. Chen H, Chen S, Chiu W, Lee L, Hung C, Hung C *et al.* A nationwide epidemiological study of spinal cord injury in geriatric patients in Taiwan. *Neuroepidemiology* 1997; **16**(5): 241-247.
6. Yang N-P, Deng C-Y, Lee Y-H, Lin C-H, Kao C-H, Chou P. The incidence and characterisation of hospitalised acute spinal trauma in Taiwan—A population-based study. *Injury Int J Care injured* 2008; **39**: 443-450.
7. Chen CF, Lien IN. Spinal Cord Injuries in Taipei, Taiwan, 1978-1981. *Paraplegia* 1985; **23**: 364-370.
8. Lan C, Lai JS, Chang KH, Jean YC, Lien IN. Traumatic spinal cord injuries in the rural region of Taiwan: an epidemiological study in Haulien county, 1986-1990. *Paraplegia* 1993; **31**: 398-403.
9. Hoque MF, Grangeon C, Reed K. Spinal cord lesions in Bangladesh: an epidemiological study 1994 -1995. *Spinal Cord* 1999; **37**: 858-861.
10. Minaire P, Castanier M, Girard R, Berard E, Deidier C, Bourret J. Epidemiology of spinal cord injury in the Rhone-Alpes Region, France, 1970-75. *Paraplegia* 1978-79; **16**: 76-87.
11. Lakhey S, Jha N, Shrestha BP, Niraula S. Aetioepidemiological profile of spinal injury patients in Eastern Nepal. *Tropical Doctor* 2005; **35**(4): 231-3.
12. Raja IA, Viobra AH, Ahmed M. Neurotrauma in Pakistan. *World J. Surgery* 2001; **25**: 1230-1237.
13. Masood Z, Wardug GM, ashraf J. Spinal Injuries: Experience of a local neurosurgical centre. *Pak J Med Sci* 2008; **24**(3): 368-371.
14. Rathore MFA, Hanif S, Farooq F, Ahmad N, Mansoor SN. Traumatic spinal cord injuries at a tertiary care rehabilitation institute in Pakistan. *JPMA - Journal of the Pakistan Medical Association* 2008; **58**(2): 53-7.

15. Kovindha A. A retrospective study of spinal cord injuries at Maharaj Nakorn Chiang Mai Hospital, during 1985-1991. *Chiang Mai Med Bull* 1993; **32**(2): 85-92.
16. Pajareya K. Traumatic spinal cord injuries in Thailand; an epidemiologic study in Siriraj Hospital, 1989-1994. *Spinal Cord* 1996; **34**: 608-610.
17. Weerts E. *Prevention of spinal cord injury accidents. What does it mean? How can I avoid it? How can I help?*
: Hanoi, 2008.
18. Weerts E. *Final reporting of project outcomes Spinal Cord Injury care and Orthopedic workshop*: Hanoi, 2009.
19. O'Connor P. Incidence and patterns of spinal cord injury in Australia. *Accident Analysis & Prevention* 2002; **34**(4): 405-15.
20. Cripps R. Spinal cord injury, Australia, 2006–07. In: *Injury Research and Statistics Series Number 48. Cat. no. INJCAT 119*. Adelaide: AIHW, 2008.
21. Cheshire DJE. *The Complete and Centralized Treatment of Paraplegia: A Report on the Spinal Injuries Centre for Victoria, Australia*. National Institute for Neurological Diseases Workshop: Melbourne, Victoria, 1966.
22. Sutton NG. *Injuries to the Spinal Cord. The management of paraplegia and tetraplegia.*, Butterworth: London, pg 185, 1973, 185pp.
23. Burke DC, Burley HT, Ungar GH. Data on spinal injuries - Part 1. Collection and analysis of 352 consecutive admissions. *Aust NZ J Surg* 1985; **55**: 3-12.
24. Dixon GS, Danesh JN, Caradoc-Davies TH. Epidemiology of spinal cord injury in New Zealand. *Neuroepidemiology* 1993; **12**: 88-95.
25. Jankowski R, Zukiel R, Nowak S, Czekanowska-Szlandrowicz R, stachowska-Tomczak B. Vertebral column and spinal cord injuries: isolated and concomitant with multiple injury. *Chir Narz Ruchu Ortop Pol* 1993; **58**(5): 353-359.
26. Soopramanien A. Epidemiology of spinal injuries in Romania. *Paraplegia* 1994; **32**: 715-722.
27. Sabre L, Linnamagi U, Derrik G, Rekand T, Asser T, Korv J. Traumatic spinal cord injuries in Estonia from 2003 to 2007. In: *ISCoS*. Florence: University of Tartu, 2009.
28. Silberstein B, Rabinovich S. Epidemiology of spinal cord injuries in Novosibirsk, Russia. *Paraplegia* 1995; **33**(6): 322-325.
29. Kondakov EN, Simonova IA, I.V. P. The epidemiology of injuries to the spine and spinal cord in Saint Petersburg. *Zhurnal Voprosy Neurokhirurgii Imeni N - N - Burdenko* 2002; **2**: 50-3.

30. Biering-Sorensen F, Pedersen V, Clausen S. Epidemiology of Spinal Cord Lesions in Denmark. *Paraplegia* 1990; **28**: 105-108.
31. Kannus P, Niemi S, Palvanen M, Parkkari J. Continuously Increasing Number and Incidence of Fall-Induced, Fracture-Associated, Spinal Cord Injuries in Elderly Persons. *Arch Intern Med* 2000; **160**(July): 2145-2149.
32. Dahlberg A, Kotila M, Leppanen P, Kautiainen H, Alaranta H. Prevalence of spinal cord injury in Helsinki. *Spinal Cord* 2005; **43**(1): 47-50.
33. Ahoniemi E, Alaranta H, Hokkinen E-M, Valtonen K, Kautiainen H. Incidence of traumatic spinal cord injuries in Finland over a 30-year period. *Spinal Cord* 2008; **46**(1): 1-4.
34. Albert T, Ravaud JF, Tetrafigap g. Rehabilitation of spinal cord injury in France: a nationwide multicentre study of incidence and regional disparities. *Spinal Cord* 2005; **43**(6): 357-65.
35. Exner G, Meinecke F-W. Trends in the treatment of patients with spinal cord lesions seen within a period of 20 years in German centers. *Spinal Cord* 1997; **35**: 415-419.
36. Koning W, Frowein RA. Incidence of Spinal Cord Injury in the Federal Republic of Germany. *Neurosurgical Review* 1989; **12**(Suppl 1): 562-566.
37. Divanoglou A, Levi R. Incidence of traumatic spinal cord injury in Thessaloniki, Greece and Stockholm, Sweden: a prospective population-based study. *Spinal Cord* 2009; **47**(1): 1-6.
38. Pedersen V, Muller PG, Biering-Sorensen F. Traumatic Spinal Cord Injuries in Greenland 1965-1986. *Paraplegia* 1989; **27**: 345-349.
39. Knutsdottir S. Spinal cord injuries in Iceland 1973-1989. A follow up study. *Paraplegia* 1993; **31**: 68-72.
40. Knutsdottir S, Thorisdottir H. Epidemiology of Spinal Cord Injuries in Iceland 1973-2008. In: *ISCoS*. Florence, Italy, 2009.
41. O'Connor RJ, Murray PC. Review of spinal cord injuries in Ireland. *Spinal Cord* 2006; **44**(7): 445-8.
42. Roche SJ, Sloane PA, McCabe JP. Epidemiology of spine trauma in an Irish regional trauma unit: A 4-year study. *Injury Int J Care injured* 2008; **39**: 436-442.
43. Catz A, Thaleisnik M, Fishel B, Ronen J, Spasser R, Fredman B *et al*. Survival following spinal cord injury in Israel. *Spinal Cord* 2002; **40**(11): 595-598.
44. Ohry A, Rozin R, Brooks ME. Paediatric traumatic Spinal cord Injuries in Israel. *Isr J med Sci* 1985; **21**: 526-528.

45. Celani MG, Spizzichino L, Ricci S, Zampolini M, Franceschini M. Spinal Cord Injury in Italy: A Multicenter Retrospective Study. *Arch Phys Med Rehabil* 2001; **82**(May): 589-596.
46. Pagliacci MC, Celani MG, Zampolini M, Spizzichino L, Franceschini M, Baratta S *et al.* An Italian Survey of Traumatic Spinal Cord Injury. The Gruppo Italiano Studio Epidemiologico Mielolesioni Study. *Arch Phys Med Rehabil* 2003; **84**(September): 1266-1275.
47. Caldana L, Lucca L. Epidemiological remarks on traumatic spinal cord injuries and non-traumatic spinal cord diseases in Veneto 1994–1995. *Eur Med Phys* 1998; **34**: 159-168.
48. Schonherr MC, Groothoff JW, Mulder GA, Eisma WH. Rehabilitation of patients with spinal cord lesions in the Netherlands: an epidemiological study. *Spinal Cord* 1996; **34**: 679-683.
49. van Asbeck F, Post M, Pangalila R. An epidemiological description of spinal cord injuries in The Netherlands in 1994. *Spinal Cord* 2000; **38**(7): 420-424.
50. Osterthun R, Post MWM, van Asbeck FWA. Characteristics, length of stay and functional outcome of patients with spinal cord injury in Dutch and Flemish rehabilitation centres. *Spinal Cord* 2009; **47**(4): 339-344.
51. Lidal IB, Snekkevik H, Aamodt G, Hjeltnes N, Stanghelle JK, Biering-Sorensen F. Mortality after Spinal Cord Injury in Norway. *J Rehabil Med* 2007; **39**: 145-151.
52. Gjone R, Nordlie L. Incidence of traumatic paraplegia and tetraplegia in Norway: A statistical survey of the years 1974 and 1975. *Paraplegia* 1978-79; **16**: 88-93.
53. Augutis M, Abel R, Levi R. Pediatric spinal cord injury in a subset of European countries. *Spinal Cord* 2006; **44**(2): 106-12.
54. Hagen EM, Eide GE, Rekand T, Gilhus NE, Gronning M. A 50-year follow-up of the incidence of traumatic spinal cord injuries in Western Norway. *Spinal Cord* 2010; **48**(4): 313-318.
55. Martins F, Freitas F, Martins L, Dartigues J, Barat M. Spinal cord injuries--epidemiology in Portugal's central region. *Spinal Cord* 1998; **36**(8): 574-578.
56. Herruzo-Cabrera R, Garcia-Reneses J, Vizcaino-Alcaide MJ, Madero R, Gil-Miguel A, Rey-Calero J. Epidemiology of traumatic spinal cord injury in Spain during 1984-1985. *Rev Clin Esp* 1993; **192**: 217-222.
57. Augutis m, Levi R. Pediatric spinal cord injury in Sweden: incidence, etiology and outcome. *Spinal Cord* 2003; **41**(6): 328-336.
58. Gehrig R. Statistics of acute paraplegia and tetraplegia on a national scale, Switzerland 1960-1967. *Paraplegia* 1988; **6**(2): 93-95.

59. Aung TS, El Masry WS. Audit of a British Centre for spinal injury. *Spinal Cord* 1997; **35**: 147-150.
60. Brasil AVB, Coelho DG. The neurological outcome of acute spinal cord injury in a neurosurgical hospital of a developing country. *Spinal Cord* 1998; **36**: 353-356.
61. Greve J. Epidemiology from the developing world: what we have, what we need -- Brazil. In. Medical School Of the University of Sao Paulo, Sao Paulo, Brazil, 2009. pp 1-8.
62. Barros F, Taricco MA, Oliveira RP, Greve JM, Santos LC, Napoli MM. Epidemiological study of patients with spinal cord injuries. *Rev hosp Fac Med S Paulo* 1990; **45**(3): 123-126.
63. Campos da Paz A, Beraldo PSS, Almeida MCRR, Neves EGC, Alves CMF, Khan P. Traumatic injury to the spinal cord. Prevalence in Brazilian hospitals. *Paraplegia* 1992; **30**: 636-640.
64. Otom A, Doughan A, Kawar J, Hattar E. Traumatic spinal cord injuries in Jordan--an epidemiological study. *Spinal Cord* 1997; **35**(4): 253-255.
65. Alshahri S. Traumatic spinal cord injury in Saudi Arabia: An epidemiological study. Unpublished work. In, 2010.
66. Dincer F, Oflazer A, Beyazova M, Celiker R, Basgoze O. Traumatic spinal cord injuries in Turkey. *Paraplegia* 1992; **30**: 641-646.
67. Karacan I, Kovuncu H, Pekel O, Sumbuloglu G, Kimap M, Dursun H *et al.* Traumatic spinal cord injuries in Turkey: a nation-wide epidemiological study. *Spinal Cord* 2000; **38**(11): 697-701.
68. Gur A, Kemaloglu MS, Cevik R, Sarac AJ, Nas K, Kapukaya A *et al.* Characteristics of traumatic spinal cord injuries in south-eastern Anatolia, Turkey: a comparative approach to 10 years' experience. *International Journal of Rehabilitation Research* 2005; **28**(1): 57-62.
69. Karamehmetoglu S, Unal S, Karacan I, Yilmaz H, Togay H, Ertekin M *et al.* Traumatic spinal cord injuries in Istanbul, Turkey. An epidemiological study. *Paraplegia* 1995; **33**(8): 469-471.
70. Karamehmetoglu S, Nas K, Karacan I, Sarac A, Koyuncu H, Ataoglu S *et al.* Traumatic spinal cord injuries in Southeast Turkey: an epidemiological study. *Spinal Cord* 1997; **35**(8): 531-533.
71. Rick Hansen Spinal Cord Injury Register. *Spinal Cord Injury Facts and Statistics: Canada*, 2006.
72. Tator CH, Duncan EG, Edmonds VE, Lapczak LI, Andrews DF. Changes in Epidemiology of Acute Spinal Cord Injury from 1947 to 1981. *Surg Neurol* 1993; **40**: 207-215.

73. Barss P, Djerrari H, Leduc BE, Lepage Y, Dionne CE. Risk factors and prevention for spinal cord injury from diving in swimming pools and natural sites in Quebec, Canada: A 44-year study. *Accident Analysis & Prevention* 2008; **40**: 787-797.
74. Andrews LG, Jung SK. Spinal cord injuries in children in British Columbia. *Paraplegia* 1979; **17**: 442-451.
75. Hamilton mG, Myles ST. Pediatric spinal injury: review of 174 hospital admissions. *J. Neurosurg* 1992; **77**: 700-704.
76. Hu R, Mustard CA, Burns C. Epidemiology of incident spinal fracture in a complete population. *Spine* 1996; **21**(4): 492-499.
77. Pickett W, Simpson K, Walker J, Brison RJ. Traumatic spinal cord injury in Ontario, Canada. *Journal of Trauma Injury, Infection, and Critical Care* 2003; **55**(6): 1070-6.
78. Dryden D, Saunders L, Rowe B, May L, Yiannakoulis N, Svenson L *et al.* The epidemiology of traumatic spinal cord injury in Alberta, Canada. *Canadian Journal of Neurological Sciences* 2003; **30**(2): 113-121.
79. Pickett GE, Campos-Benitez M, Keller JL, Duggal N. Epidemiology of traumatic spinal cord injury in Canada. *Spine* 2006; **31**(7): 799-805.
80. Samsa GP, Patrick CH, Feussner jR. Long-term Survival of Veterans with Traumatic Spinal Cord Injury. *Arch Neurol* 1993; **50**: 909-914.
81. Bracken MB, Freeman J, D.H., Hellenbrand K. Incidence of Acute Traumatic Hospitalized Spinal Cord Injury in the United States, 1970-1977. *American Journal of Epidemiology* 1981; **113**(6): 615-622.
82. Nobunaga AI, Go BK, Karunas RB. Recent demographic and Injury Trends in People Served by the Model Spinal Cord Injury Care systems. *Arch Phys Med Rehabil* 1999; **80**(November): 1372-1382.
83. Jackson AB, Dijkers M, DeVivo MJ, Poczatek RB. A Demographic Profile of New Traumatic Spinal Cord Injuries: Change and Stability Over 30 Years. *Arch Phys Med Rehabil* 2004; **85**(November): 1740-1748.
84. Kalsbeek WD, McLaurin RL, Harris 111 BSH, Miller JD. The National Head and Spinal Cord Injury Survey: Major Findings. *J. Neurosurg.* 1980; **53**: 519-531.
85. Burney RE, Maio RF, Maynard F, Karunas RB. Incidence, characteristics, and outcome of spinal cord injury at trauma centers in North America. *Arch Surg* 1992; **128**: 596-599.
86. Stover SL, Fine PR. The epidemiology and economics of spinal cord injury. *Paraplegia* 1987; **25**: 225-228.

87. Harvey C, Wilson SE, Greene CG, Berkowitz M, Stripling TE. New estimates of the direct costs of traumatic spinal cord injuries: results of a nationwide survey. *Paraplegia* 1992; **30**: 834-850.
88. Ditunno JF, Formal CS. Chronic spinal cord injury. *New Engl J Med* 1994; **330**: 550-556.
89. Lasfargues JE, Custis D, Morrone F, Carswell J, Nguyen T. A model for estimating spinal cord injury prevalence in the United States. *Paraplegia* 1995; **33**(2): 62-8.
90. Vitale MG, Goss JM, Matsumoto H, Roye DP, Jr. Epidemiology of pediatric spinal cord injury in the United States: years 1997 and 2000. *J Pediatr Orthop* 2006; **26**(6): 745-9.
91. Ho CH, Wuermser L-A, Priebe MM, Chiodo AE, Scelza WM, Kirshblum SC. Spinal cord injury medicine. 1. Epidemiology and classification. *Archives of Physical Medicine & Rehabilitation* 2007; **88**(3 Suppl 1): S49-54.
92. National Spinal Cord Injury Statistical Center. Spinal cord injury. Facts and figures at a glance. *Journal of Spinal Cord Medicine* 2005; **28**(4): 379-80.
93. Griffin MR, Opitz JL, Kurland LT, Ebersold MJ, O'Fallon WM. Traumatic Spinal Cord Injury in Olmstead County, Minnesota, 1935-1981. *American Journal of Epidemiology* 1985; **121**(6): 884-895.
94. Kraus JF, Franti CE, Riggins RS, Richards D, Borhani NO. Incidence of Traumatic spinal Cord Lesions. *J Chron Dis* 1975; **28**: 471-492.
95. Acton PA, Farley T, Freni LW, Ilegbodun VA, Sniezek JE, Wohlleb JC. Traumatic spinal cord injury in Arkansas, 1980-1989. *Arch Phys Med Rehabil* 1993; **74**: 1035-1040.
96. Morbidity and Mortality Weekly Report. Trends in traumatic spinal cord injury - New York 1982-1988. 1991; **40**(31): 535-537.
97. Goebert DA, Ng MY, Varney JM, Sheetz DA. Traumatic spinal cord injury in Hawaii. *Hawaii Medical Journal* 1991; **50**(2): 44-50.
98. Price C, Makintubee S, Herndon W, Istre GR. Epidemiology of Traumatic Spinal Cord Injury and Acute Hospitalization and Rehabilitation Charges for Spinal Cord Injuries in Oklahoma, 1988-1990. *American Journal of Epidemiology* 1994; **139**(1): 37-47.
99. Thurman DJ, Burnett CL, Jeppson L, Beaudoin DE, Sniezek JE. Surveillance of spinal cord injuries in Utah, USA. *Paraplegia* 1994; **32**: 665-669.
100. Warren S, Moore M, Johnson M. Traumatic head and spinal cord injuries in Alaska (1991-1993). *Alaska Medicine* 1995; **37**(1): 11-19.

101. Surkin J, Gilbert B, Harkley H, Sniezek J, Currier M. Spinal cord injury in Mississippi. Findings and evaluation, 1992-1994. *Spine* 2000; **25**(6): 716-721.
102. Johnson RL, Gabella BA, Gerhart KA, McCray J, Menconi jC, whiteneck GG. Evaluating Sources of Traumatic Spinal Cord Injury Surveillance Data in Colorado. *American Journal of Epidemiology* 1997; **146**(3): 266-272.
103. Burke D, Linden R, Zhang Y, Maiste A, Shields C. Incidence rates and populations at risk for spinal cord injury: A regional study. *Spinal Cord* 2001; **39**(5): 274-278.
104. Buechner JS, Speare MC, Fontes J. Hospitalizations for Spinal Cord Injuries, 1994-1998. *Medicine and Health/Rhode Island* 2000; **83**(3): 92-93.
105. Calancie B, Molano MR, Broton JG. Epidemiology and demography of acute spinal cord injury in a large urban setting. *Journal of Spinal Cord Medicine* 2005; **28**(2): 92-6.
106. Maharaj J. Epidemiology of spinal cord paralysis in Fiji: 1985-1994. *SpinalCord* 1996; **34**(9): 549-559.
107. Key AG, Retief PjM. Spinal Cord Injuries: An Analysis of 300 New Lesions. In: *International Medical Society of Paraplegia: Annual Scientific Meeting*. Tel-Aviv: Paraplegia, 1968. pp 243-249.
108. Velmahos GC, Degiannis E, Hart K, Souter I, Saadia R. Changing Profiles in Spinal Cord Injuries and Risk Factors Influencing Recovery after Penetrating Injuries. *The Journal of Trauma: Injury, Infection, and Critical Care* 1995; **38**(3): 334-337.
109. Hart C, Williams E. Epidemiology of spinal cord injuries: a reflection of changes in South African society. *Paraplegia* 1994; **32**: 709-714.
110. Levy LF, Makarawo S, Madzivire D, Bhebhe E, Verbeek N, Parry O. Problems, struggles and some success with spinal cord injury in Zimbabwe. *Spinal Cord* 1998; **36**: 213-218.
111. Iwegbu CG. traumatic paraplegia in Zaria, Nigeria: The case for a centre for injuries of the spine. *Paraplegia* 1983; **21**: 81-85.
112. Igun GO, Obekpa OP, Ugwu BT, Nwadiaro HC. Spinal injuries in the Plateau State, Nigeria. *East African Medical Journal* 1999; **76**(2): 75-79.
113. Obalum DC, Giwa SO, Adekoya-Cole TO, Enweluzo GO. Profile of spinal injuries in Lagos, Nigeria. *Spinal Cord* 2009; **47**(2): 134-137.
114. Solagberu BA. Spinal cord injuries in Ilorin, Nigeria. *West Afr J Med* 2002; **21**(3): 230-232.
115. Nwankwo OE, Katchy AU. Outcome of a 12-week programme for management of the spinal cord injured with participation of patient's relations at Hilltop Orthopaedic Hospital, Enugu, Nigeria. *Spinal Cord* 2003; **41**(2): 129-133.

116. Nwadinigwe CU, Iloabuchi TC, Nwabude IA. Traumatic Spinal Cord Injuries (SCI): A Study of 104 Cases. *Nigerian Journal of Medicine* 2004; **13**(2): 161-165.
117. Olasode BJ, Komolafe IE, Komolafe M, Olasode OA. Traumatic spinal cord injuries in Ile-Ife, Nigeria, and its environs. *Tropical Doctor* 2006; **36**(3): 181-2.
118. Gosselin RA, Coppotelli C. A follow-up study of patients with spinal cord injury in Sierra Leone. *international Orthopaedics (SICOT)* 2005; **29**: 330-332.